

A Qubit on the Faraday Mirror

F. De Martini, F. Sciarrino and V. Secondi

Dipartimento di Fisica and

Istituto Nazionale per la Fisica della Materia

Università di Roma "La Sapienza", Roma, 00185 - Italy

Abstract

In the context of Quantum Information (QI) the "Faraday Mirror" acts as a *non-universal* NOT Gate. As such its behaviour complies with the principles of quantum mechanics. This non trivial result, at the core of some recent misinterpretations in the QI community, has been reached by a thorough experimental investigation of the properties of the device including the adoption of modern *Quantum Process Tomography*. In addition, the "universal optical compensation" method devised by Mario Martinelli, of common use in long distance quantum-cryptography, has been fully investigated theoretically and experimentally.

The today common interest in the "optimal" realization of "forbidden" quantum information processes and the recent realizations of the Universal Not Gate [1–3] have stressed the somewhat intriguing role of the "Faraday Mirror" (*FRM*) in connection with the "spin flipping" process, i.e. the *positive* map (P-map) that should transform any qubit $|\Psi\rangle$ into the orthogonal one $|\Psi_\perp\rangle$, i.e. such as $\langle\Psi|\Psi_\perp\rangle = 0$ [4]. In other words, this map should draw any point P representative of $|\Psi\rangle$ on the Poincaré sphere into its "antipode" P' representative of $|\Psi_\perp\rangle$. In this connection the *FRM* issue has been taken up by an extended comment in a recent review paper on Quantum Cryptography [5]. In view of some recent misinterpretations of the phenomenon we do believe that this matter requires a necessary clarification [5].

The *FRM* is a simple, somewhat exotic device consisting of an optical mirror (M) that reflects a monochromatic light beam, associated with a (single gaussian) mode $\mathbf{k}$, into the mode $-\mathbf{k}$. In front of the mirror is placed a "Faraday Rotator" (*FR*), a laboratory device generally used to avoid spurious laser back-reflection effects, in which a transparent paramagnetic spin glass rod is acted upon by a static magnetic field $\vec{B}$, parallel to $\mathbf{k}$. By the magneto-optical Faraday effect the incoming polarization (π) is "rotated" by keeping the same π -character: a "linear" π is transformed into a "linear" π' , a "circular" π into a "circular" π' etc. In particular, the field $\vec{B}$ and the π -rotation "angle" θ can be "tuned" as to realize the orthogonality between π and π' [6]. Indeed, precisely for this useful property the *FRM* device is commercially advertized in optics and in laser technology. Furthermore, according to an original proposal by Mario Martinelli [7], it can also be adopted to eliminate spurious birefringence effects in fiber optics technology, as we shall see.

From a quantum mechanical perspective this process, as it is commonly understood, may indeed represent a real puzzle. The obvious point of concern is precisely his alleged property of realizing a *universal* spin-flip transformation $|\Psi\rangle \rightarrow |\Psi_\perp\rangle$ in spite of this map being "forbidden" by quantum mechanics (*QM*). Indeed according to *QM* only the *complete-positive* maps (CP-maps) are realizable by Nature. In addition, since the *FRM* is a *passive*, and then noiseless device, that transformation should be *exact*, i.e. implying a *full transfer*

of quantum information from $|\Psi\rangle$ to $|\Psi_\perp\rangle$. In order to avoid further speculations, the present paper intends to clarify the problem by a clear-cut recourse to the experiment.

A Spontaneous Parametric Down Conversion (SPDC) process was excited in a 1.5 mm thick BBO (β -barium borate) nonlinear (NL) crystal slab, by a mode-locked pulsed UV beam with wavelength (wl) λ_p : Figure 1. By this setup entangled photon pairs in a *singlet-state* of polarization (π) were created with common wl's $\lambda = \frac{1}{2}\lambda_p = 795nm$. Each excitation laser pulse had a time duration $\tau \approx 140$ fsec. Because of the low pump intensity, the probability of the unwanted simultaneous $N = 2$ pairs generation in the NL crystal was found $\approx 10^{-2}$ smaller than the one for a single pair: $N = 1$. In virtue of the nonlocal correlation existing between the photons of each pair, the qubit $|\Psi\rangle$ injected over the mode $\mathbf{k}_2$ into the *FRM* device was deterministically selected by the measurement apparatus coupled to mode $\mathbf{k}_1$. This one consisted of a π -analyzer, i.e. a polarizing beam-splitter (*PBS*), a pair of $(\lambda/2 + \lambda/4)$ optical wave-plates (wpl) and of a tunable Babinet Compensator (*BC*). The detector D_1 , and the end detector D_2 , were equal SPCM-AQR14 Si-avalanche single photon units with quantum efficiencies $QE's \cong 0.55$. One interference filter with bandwidth $\Delta\lambda = 3nm$ was placed in front of each D . The Faraday rotator (*FR*), a CVI-FR-660/1100-8 device accurately $\vec{B}$ -tuned for the wl λ , could be optionally removed from the optical network. It was terminated by a 100% reflectivity mirror M . In order to avoid the use of other mirrors, the output qubit $|\Psi'\rangle$ emitted by *FRM* over the mode $-\mathbf{k}_2$ was transmitted through the transparent BBO NL slab and finally detected by the end measurement apparatus, made of a *PBS*, a pair of $(\lambda/2 + \lambda/4)$ wpl set and a detector D_2 . Three quartz slabs (Q) inserted on the modes $\mathbf{k}_1$, $\mathbf{k}_2$, $-\mathbf{k}_2$ and the Babinet compensator on mode $\mathbf{k}_1$ secured the accurate compensation of all residual birefringence effects, e.g. of the BBO slab.

In a first experiment we wanted to fully characterize by standard quantum tomography the input entangled state generated by SPDC, *in absence* of *FRM* [8]. The reconstructed density matrix ρ_{k_1, k_2}^{in} of the photon pair is reported in Fig. 2-a. There the realization of the *singlet* input pure state condition $|\Phi\rangle = 2^{-1/2} (|H\rangle_1 |V\rangle_2 - |V\rangle_1 |H\rangle_2)$ is clearly shown, where $|H\rangle_i$ and $|V\rangle_i$ stands for *linear- π* horizontal and vertical qubits, associated respectively to

modes $\mathbf{k}_i$, $i = 1, 2$. We then analyzed the transformation induced by the mirror reflection only (a condition henceforth referred to as: *M-reflection*) and by the complete *FRM* device (*FRM-reflection*). By the knowledge of $|\Phi\rangle$ we prepared the input qubit $|\Psi\rangle$ injected over $\mathbf{k}_2$ by exploiting the nonlocal π -correlation existing between the two photons of each pair. Precisely, we limited ourselves to prepare six input qubits $|\Psi\rangle : |H\rangle, |V\rangle, |L_{\pm}\rangle \equiv 2^{-\frac{1}{2}}(|H\rangle \pm |V\rangle), |C_{\pm}\rangle \equiv 2^{-\frac{1}{2}}(|H\rangle \pm i|V\rangle)$. The results of two experiments carried out respectively in absence and in presence of *FR*, were the following:

M-reflection: $|H\rangle \rightarrow |H\rangle; |V\rangle \rightarrow |V\rangle; |L_{\pm}\rangle \rightarrow |L_{\mp}\rangle; |C_{\pm}\rangle \rightarrow |C_{\mp}\rangle$, with fidelities $\mathcal{F} = 0.98, 0.98, 0.90, 0.90$ respectively.

FRM-reflection: $|H\rangle \rightarrow |V\rangle; |V\rangle \rightarrow |H\rangle; |L_{\pm}\rangle \rightarrow |L_{\pm}\rangle; |C_{\pm}\rangle \rightarrow |C_{\mp}\rangle$ with $\mathcal{F} = 0.99, 0.99, 0.92, 0.92$ respectively.

Here the arrows indicate the mapping from the input to the output states, associated with the input/output modes $\mathbf{k}_2$ and $-\mathbf{k}_2$, correspondingly. The $\mathcal{F}$ values, here defined as the statistical fractions of the "right" outcomes, were limited by an imperfect correction of the optical walk-off inside the NL crystal. We can see that, in agreement with *QM*, the *FRM* device *does not* realize the *universal* spin-flip action but rather a *CP-map*, i.e. a *unitary* transformation expressible by a definite rotation around the center of the Poincare' sphere of the unit vector $\vec{p}$. This vector is then drawn by the transformation into a position $\vec{p}'$ in the tri-dimensional *spin space* with orthogonal axes $\sigma_1, \sigma_2, \sigma_3$: Figure 1, inset. Correspondingly, the vertex of $\vec{p}$, i.e. the point P expressing the input $|\Psi\rangle$ on the surface of the Poincare' sphere is drawn into the point P' expressing the output $|\Psi'\rangle$.

Let's gain insight into the overall process by following the path through *FRM* of the short light pulse associated with any single photon carrying the qubit. Consider the (real) spatial reference frame (x,y,z) the qubit is referred to before reflection by *M*: let the axes x, y belong to the plane of *M*, while the z-axis, orthogonal to *M*, is parallel to and oriented as the input photon momentum $\hbar\mathbf{k}_2$. Assume that the qubit's π -states are identified according to the *right-hand* (r.h.) coordinate convention, henceforth referred to as "*frame-convention*". Of course, the preservation of the *frame-convention* upon *M-* or *FRM-reflection* is a

necessary requirement for a correct understanding of the reflection dynamics, since the input and output qubits are identified precisely on the basis of the real-space reference frame. This implies that, after reflection the photon's π -states of any photon are identified on the basis of a new coordinate frame ($x'=x$, $y'=-y$, $z'=-z$), obtained by rotating the old frame around the $\vec{x}$ axis by an angle 180° . This easily explains the (apparent) puzzling behaviour with respect to state-orthogonality of the input/output *linear polarization* states under *FRM-reflection*.

In order to visualize the process with the help of the Poincare' sphere, let us consider the *M-reflection* first. Let the point P to represents the input qubit when the input photon hits M , as said: $|\Psi\rangle = \psi_1 |H\rangle + \psi_2 |V\rangle$, $|\psi_1|^2 + |\psi_2|^2 = 1$, being ψ_i the complex components of the spinor $\begin{bmatrix} \psi_1 \\ \psi_2 \end{bmatrix}$. Classical optics and our experimental results above show that the reversal of the particle motion and the preservation of the *frame-convention* upon *M-reflection* implies a rotation of $\vec{p}$ around the spin reference axis σ_3 by an angle 180° , i.e. by the unitary map: $\mathbb{U}_3 = \exp(i\frac{1}{2}\pi\sigma_3) = i\sigma_3$. Note that, after the corresponding representative point P being drawn into P' by the *M-reflection* transformation, the *same* Poincare' sphere can still be adopted for the future qubit dynamics, indeed a useful condition when searching for a possible "antipode" of P over the sphere. In summary, the *M-reflection* process transforms the input qubit $|\Psi\rangle$ above as follows: $\mathbb{U}_3 \begin{bmatrix} \psi_1 \\ \psi_2 \end{bmatrix} = i \begin{bmatrix} \psi_1 \\ -\psi_2 \end{bmatrix}$ [9].

Let us turn now our attention to the *FRM-reflection*. If P now represents the qubit before entering the device, the first passage through *FR* draws P into O by a positive rotation by 90° around σ_2 , i.e. corresponding to the evolution operator: $\mathbb{U}_{2+} = \exp(i\frac{1}{4}\pi\sigma_2)$. The *M-reflection* contributes with the map $\mathbb{U}_3$, leading from the point O to O' , as just seen. The second passage through *FR* after *M-reflection* is represented by $\mathbb{U}_{2-} = \exp(-i\frac{1}{4}\pi\sigma_2)$ since the change of the spatial reference frame leaves unchanged the (axial) magnetic-polarization pseudo-vector $\vec{\mathcal{M}} \propto \vec{B}$ while the sign of the phase $\pi/2$ is inverted because of the particle's propagation reversal [10]. In summary, the overall evolution of the input qubit $|\Psi\rangle$ through the *FRM* leaves it in the final state: $|\Psi'\rangle = \mathbb{U}_{FRM} |\Psi\rangle = \mathbb{U}_{2-} \mathbb{U}_3 \mathbb{U}_{2+} |\Psi\rangle$

$= \frac{1}{2}(1 - i\sigma_2) \times (i\sigma_3) \times (1 + i\sigma_2) |\Psi\rangle = i\sigma_1 |\Psi\rangle$, since: $\sigma_i\sigma_j = i\sigma_k$ for an even permutation of i, j, k . Correspondingly, the initial point P on the sphere is drawn into the successive positions O , O' and P' . Correspondingly, the overall *FRM* process results in a overall 180° rotation around the axis σ_1 of the initial representative vector $\vec{p}$, i.e. the input qubit is acted upon by the *unitary*: $\mathbb{U}_{FRM} = \exp(i\frac{1}{2}\pi\sigma_1) = i\sigma_1$, as seen. As we can see, the final representative point P' is *not* the antipode of *any possible* P on the sphere and then *FRM does not* realize the *universal* quantum-NOT Gate, as expected from QM. Indeed the spin-flip action is realized exactly *only* for all representative points P belonging to the equatorial plane orthogonal to the σ_1 axis, such as for the qubits $|H\rangle$, $|V\rangle$ and $|C_\pm\rangle$.

As an additional deeper-level demonstration, a very general characterization of the *FRM* device was attained experimentally by the entanglement-assisted *Quantum Process Tomography* (*QPT*) [11]. Let us outline the basics of this approach, by first representing the state of the input qubit corresponding to a point P on the Poincare' sphere, by the density operator $\rho_{k2} = \frac{1}{2}(\mathbb{I} + \vec{r} \cdot \vec{\sigma})$. Then this qubit can be represented by a 4-dimensional vector $(1, \vec{r})$. The single map $\mathcal{E}(\rho)$, expressing the *FRM-reflection*, is completely described by a 4×4 real matrix $\mathbf{M}_{FRM}$, which maps ρ_{k2} into the density matrix: $\rho'_{-k2} = \mathbf{M}_{FRM}\rho_{k2}$ [12]. A most important, exclusive property of the *QPT* method is the exploitation of the *quantum parallelism* associated with the entanglement condition [11]. Indeed, in our experiment the matrix $\mathbf{M}_{FRM}$ was experimentally reconstructed with the use of the input pure *entangled state* $\rho_{k1,k2}^{in}$ of the two correlated qubits. The first step consisted of performing the quantum state tomography of that state $\rho_{k1,-k2}^{in}$ (Fig.2-a-left side). Then by *QPT* the qubit associated with mode $\mathbf{k}_1$, call it "qubit 1", was left unchanged while the "qubit 2", associated with the mode $\mathbf{k}_2$ underwent the $\mathcal{E}(\rho)$ -transformation. Note that this procedure implies the investigation of the unknown map $\mathcal{E}(\rho)$, i.e. of *FRM-reflection*, by a *complete* span over the Hilbert space $\mathcal{H}_2$ of the injected "qubit 2" because of its *mixed-state* condition. The final state of the two qubits $\rho_{k1,-k2}^{out} = \mathbb{I}_{k1} \otimes \mathcal{E}_{k2}(\rho_{k1,k2}^{in})$ was again investigated by tomography (Fig.2-a-right side). Finally, the matrix $\mathbf{M}_{FRM}$ was estimated by means of the experimen-

tally determined density matrices $\rho_{k1,k2}^{in}$ and $\rho_{k1,-k2}^{out}$ by adoption of an inversion software [11]. Remind the result $\mathbb{U}_{FRM} = i\sigma_1$ obtained by the theoretical analysis of the *FRM-reflection*, given above. We may then compare the matrix $\mathbf{M}_{FRM}$ associated to the unknown *FRM* map $\mathcal{E}(\rho)$ with the corresponding one expressing the σ_1 Pauli operator:

$$\mathbf{M}_{\sigma_1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \quad (1)$$

The overlap of two generic maps $\mathcal{E}$ and $\mathcal{L}$ is usually defined by the fidelity: $\mathcal{F}(\mathcal{E}, \mathcal{L}) = \int d\Psi \mathcal{F}[\mathcal{E}(|\Psi\rangle\langle\Psi|), \mathcal{L}(|\Psi\rangle\langle\Psi|)]$ [13]. In the present context we obtain: $\mathcal{F}(\sigma_1, \mathcal{E}(\rho)) = \frac{1}{2} + \frac{1}{3}(M_{22} - M_{33} - M_{44}) = (99.7 \pm 0.3)\%$ where M_{ii} is the ii -element of the matrix $\mathbf{M}_{FRM}$. The *QPT* reconstruction of $\mathbf{M}_{FRM}$ is shown in Figure 2-b). We may compare the diagram reported there with the structure of the matrix $\mathbf{M}_{\sigma_1}$, Eq.1. The correspondence is quite impressive [14].

In summary, our experimental results support the previous theoretical analysis and then the *FRM* "puzzle" can be considered fully clarified. In addition, these conclusions emphasize once again the relevance of the two different *Universal-Not Gates* ($U - NOT$) schemes that have been recently reported in the literature [2,3]. Indeed these schemes fully implement the *universality* condition, albeit *optimally*, i.e. by a fidelity $F < 1$. This last condition is physically achieved in the *deterministic, parametric amplification* $U - NOT$ scheme by addition to the output "signal" of a "noise" contribution arising from the QED vacuum fluctuations [2]. In the *probabilistic*, linear *Tele-U - NOT* scheme, a partial cancellation of information is due to the statistical, i.e. non deterministic character of the device [3].

For the sake of completeness, let's investigate here also the interesting "*FRM* - compensation" process discovered by Mario Martinelli that has recently found a widespread application whenever an active compensation of optical path fluctuations is required, as in long distance quantum cryptography applications with optical fibers [5,7]. The optical

scheme of the process is the following. Suppose that Alice (A) and Bob (B), two receiving/sending QI stations are connected by an optical device (*PC* in Figure 1) that acts on the transmitted optical signal by an (presumably unwanted) *unitary* transformation $\mathbb{U}$, e.g. corresponding to slowly fluctuating birefringence effects due possibly to temperature changes or to local changes of the curvature of the optical fiber etc. If the overall communication process implies the transmission of the signal from A to B and then back from B to A, the spurious $\mathbb{U}$ effect can be cancelled if the corresponding device (e.g. the optical fiber) is terminated by a *FRM* device. For instance, if the *FRM* device is placed at the site B the signal sent along the fiber from A to B and the received back results to A to be unaffected by the disturbance introduced by the *PC* device. In summary, looking through the fiber A can only detect the effect of *FRM* and *not* of *PC*. All this can be easily analyzed with the help of the discussion above. Let the fluctuating effect contributed by *PC* and affecting the transmission from A to B be represented generally by the *unitary* $\mathbb{U} = \exp[i\frac{1}{2}(\vec{\sigma} \bullet \vec{n})\theta]$ where $\vec{n}$ is a unit vector pointing in *any* direction in the spin space. The propagation back from B to A will be then represented by a corresponding unitary $\mathbb{U}' = \exp[-i\frac{1}{2}(\vec{\sigma} \bullet \vec{n}')\theta]$ where the phase $(-\theta)$ accounts for the propagation reversal and the components of the unit vector $\vec{n}'$ in the *spin space* are connected to $\vec{n}$ as follows: $n'_1 = n_1$, $n'_2 = -n_2$, $n'_3 = -n_3$. In other words, by retracing the propagation back through the fiber the rotation-axis involved in the $\mathbb{U}'$ transformation has undergone a rotation by an angle 180° around the axis σ_1 respect to the original direction: $\vec{n} \rightarrow \vec{n}'$. All this can be explained by considering that the fluctuating optical disturbance $\mathbb{U}$, e.g. physically due to birefringence etc., consists of an electrical-polarization $\vec{\mathcal{P}}$ effect. Since $\vec{\mathcal{P}}$ is a polar ("true") vector, it is not insensitive to any *sign change* of the spatial coordinates in *real* space, e.g. the one implied by the *FRM-reflection* [10]. Furthermore, we have already shown that the input/output dynamics due to the *FRM-reflection* implies precisely a rotation by an angle 180° around σ_1 of any unit vector representative of the input/output qubits: $\vec{p} \rightarrow \vec{p}'$. The overall process can be now formally expressed by a straightforward application of spin algebra leading to the transformation affecting any qubit forwarded by A in a round trip journey through the

whole device ($PC+FRM$): $\mathbb{U}'\mathbb{U}_{FRM}\mathbb{U} \equiv \exp[-i\frac{1}{2}(\vec{\sigma} \bullet \vec{n}')\theta] \times (i\sigma_1) \times \exp[i\frac{1}{2}(\vec{\sigma} \bullet \vec{n})\theta] = i\sigma_1 = \mathbb{U}_{FRM}$, for *any* vector $\vec{n}$ and angle θ [15]. In other words, the effect of *any* possible round-trip *unitary* $\mathbb{U}$ -process, e.g. introduced by any optical fiber or by any other optical device, is exactly cancelled, i.e. "compensated", by this very clever *universal FRM* technique.

The compensation effect has been experimentally checked by an "ergodic procedure", i.e. by insertion of a stochastically time-varying unitary operation $\mathbb{U}(t)$ into the mode $\mathbf{k}_2$ between the NL crystal and the *FRM*. More precisely this device, represented by *PC* in Figure 1, and apt to simulate the most general propagation inside an optical fiber, consisted of a cascade of two equal *Pockels Cells* (*Shangai Institute of Ceramics*, risetime ≈ 1 nsec) mutually rotated by 45° and driven by a stochastic time-sequence of high-voltage electric pulses. The output density matrices $\tilde{\rho}_{k1,-k2}^{out}$ and $\rho_{k1,-k2}^{out}$ representing the states of each photon pair after propagation into the $PC+FRM$ device, respectively upon activation and inactivation of the Pockels Cells, were reconstructed by a tomographic technique. The value of the "fidelity" $\mathcal{F}(\rho_{k1,-k2}^{out}, \tilde{\rho}_{k1,-k2}^{out}) = \left(Tr \sqrt{\sqrt{\tilde{\rho}_{k1,-k2}^{out}} \rho_{k1,-k2}^{out} \sqrt{\tilde{\rho}_{k1,-k2}^{out}}}\right)^2$ expressing the overlapping of the two experimentally determined quantum states has been found: $\mathcal{F} = (99.9 \pm 0.1)\%$. This figure shows the very high degree of compensation attainable by the method.

We are indebted with Vladimir Buzek for stimulating and lively discussions on *FRM-reflection* and Marco Ricci for help with the tomographic techniques. This work has been supported by the FET European Network on Quantum Information and Communication (Contract IST-2000-29681: ATESIT), by MIUR Cofinanziamento-2002 and by PRA-INFN 2002 (CLON).

REFERENCES


- [1] N. Gisin and S. Popescu, *Phys. Rev. Lett.* **83**, 432 (1999); V. Buzek, M. Hillery and R.F. Werner, *Phys. Rev. A* **60**, R2626 (1999).
- [2] F. De Martini, V. Buzek, F. Sciarrino and C. Sias, *Nature* **419**, 815 (2002); D. Pelliccia *et al.* *Phys. Rev. A* **68**, 042306 (2003); F. De Martini, D. Pelliccia, and F. Sciarrino, *quant-ph/0312124*.
- [3] M. Ricci, F. Sciarrino, C. Sias and F. De Martini, *quant-ph/0304070* (2003) and *Physical Review Letters* (to appear). This "Tele U-NOT" scheme realizes a new quantum information protocol: the "Teleportation of a quantum-gate".
- [4] V. Buzek and N. Gisin (private communications).
- [5] N. Gisin, G. Ribordy, W. Tittel and H. Zbinden, *Rev. Mod. Phys.* **74**, 145 (2002).
- [6] Landau-Lifshitz, *Electrodynamics of Continuous Media* (Pergamon Press, New York 1960) Ch.4.
- [7] M. Martinelli, *J. Mod. Optics* **39**, 451 (1992) and: *Opt. Comm.* **72**, 341 (1989).
- [8] A.G. White, D.F.V. James, P.H. Eberhard, and P.G. Kwiat, *Phys. Rev. Lett.* **83**, 3103 (1999).
- [9] This result is at variance with a corresponding one found in Ref. 5, page 171. There the incorrect application of the *frame-convention* to the analysis of the *M-reflection* process leads to a quite unphysical *non-unitary* time-reversal transformation.
- [10] J.D. Jackson, *Classical Electrodynamics* (Wiley, New York 1975) Ch.6.
- [11] F. De Martini, A. Mazzei, M. Ricci, G.M. D'Ariano, *Phys. Rev. A* **67**, 062307 (2003); G.M. D'Ariano and P. Lo Presti, *Phys. Rev. Lett.* **86**, 4195 (2001); M. Ricci: *private communication*.
- [12] D.K.L. Oi, *quant-ph/0106035*.

- [13] M.D. Bowdrey *et al.*, *Phys. Lett. A* **294**, 258 (2002).
- [14] G.M. D'Ariano, and P. Lo Presti, *Phys. Rev. Lett.* **91**, 047902 (2003); J.B. Altepeter, *et al.*, *Phys. Rev. Lett.* **90**, 193601 (2003).
- [15] J.J. Sakurai, *Modern Quantum Mechanics* (Addison Wesley, New York, 1994) Ch.3.


Figure Captions

Figure.1. Experimental setup adopted for the characterization of the *FRM-reflection* process. INSET: Representation of the *FRM* action on the Poincare' sphere. The dotted lines (blue) represent the three transformations due to the round-trip in the Faraday rotator (Paths PO and $O'P'$) and to the mirror reflection (Path OO'). The dashed line (red) represents the overall operation due to *FRM-reflection*, i.e. a rotation around the σ_1 axis by an angle 180° .


Figure.2. (a) Tomographic representation of the input and output entangled states $\rho_{k1,k2}^{in}$ and $\rho_{k1,-k2}^{out}$. Only the *real* parts is reported here. The maximum size of the experimentally determined *imaginary* parts is respectively 0.02 and 0.05 in the corresponding reported scales. The upper left diagram has been obtained by direct measurements taken on the mode $\mathbf{k}_2$, i.e. by replacing *FRM* with the apparatus terminated by D_2 : figure 1. (b) Reconstruction by *Quantum Process Tomography* of the matrix $\mathbf{M}_{FRM}$ representing the *FRM-reflection*.



a)




ρ^{IN}



ρ^{OUT}

b)



$\mathbf{M}_{\text{FRM}}$